

J Neuroimmune Pharmacol (2009) 4:190–199  
 DOI 10.1007/s11481-009-9152-8

## ORIGINAL ARTICLE

# Increased Accumulation of Intraneuronal Amyloid $\beta$ in HIV-Infected Patients

Cristian L. Achim · Anthony Adame ·  
 Wilmar Dumaop · Ian P. Everall · Eliezer Masliah ·  
 HNRC

Received: 15 January 2009 / Accepted: 25 February 2009 / Published online: 17 March 2009  
 © The Author(s) 2009. This article is published with open access at [Springerlink.com](http://Springerlink.com)

**Abstract** In recent years, human immunodeficiency virus (HIV)-infected patients under highly active anti-retroviral therapy (HAART) regimens have shown a markedly improved general clinical status; however, the prevalence of mild cognitive disorders has increased. We propose that increased longevity with HIV-mediated chronic inflammation combined with the secondary effects of HAART may increase the risk of early brain aging as shown by intraneuronal accumulation of abnormal protein aggregates like amyloid  $\beta$  ( $A\beta$ ), which might participate in worsening the neurodegenerative process and cognitive impairment in older patients with HIV. For this purpose, levels and distribution of  $A\beta$  immunoreactivity were analyzed in the frontal cortex of 43 patients with HIV (ages 38–60) and HIV– age-matched controls. Subcellular localization of the  $A\beta$ -immunoreactive material was analyzed by double labeling and confocal

microscopy and by immunono-electron microscopy (EM). Compared to HIV– cases, in HIV+ cases, there was abundant intracellular  $A\beta$  immunostaining in pyramidal neurons and along axonal tracts. Cases with HIV encephalitis (HIVE) had higher levels of intraneuronal  $A\beta$  immunoreactivity compared to HIV+ cases with no HIVE. Moreover, levels of intracellular  $A\beta$  correlated with age in the group with HIVE. Double-labeling analysis showed that the  $A\beta$ -immunoreactive granules in the neurons co-localized with lysosomal markers such as cathepsin-D and LC3. Ultrastructural analysis by immuno-EM has confirmed that in these cases, intracellular  $A\beta$  was often found in structures displaying morphology similar to autophagosomes. These findings suggest that long-term survival with HIV might interfere with clearance of proteins such as  $A\beta$  and worsen neuronal damage and cognitive impairment in this population.

The San Diego HIV Neurobehavioral Research Center (HNRC) group is affiliated with the University of California, San Diego, the Naval Hospital, San Diego, and the Veterans Affairs San Diego Healthcare System, and includes: Director: Igor Grant, M.D.; Co-Directors: J. Hampton Atkinson, M.D., Ronald J. Ellis, M.D., Ph.D., and J. Allen McCutchan, M.D.; Center Manager: Thomas D. Marcotte, Ph.D.; Melanie Sherman; Naval Hospital San Diego: Braden R. Hale, M.D., M.P.H. (P.I.); Neuromedical Component: Ronald J. Ellis, M.D., Ph.D. (P.I.), J. Allen McCutchan, M.D., Scott Letendre, M.D., Edmund Capparelli, Pharm.D., Rachel Schrier, Ph.D.; Jennifer Marquie-Beck; Terry Alexander, R.N.; Janis Durelle; Neurobehavioral Component: Robert K. Heaton, Ph.D. (P.I.), Mariana Cherner, Ph.D., Steven Paul Woods, Psy.D., David J. Moore, Ph.D.; Matthew Dawson; Neuroimaging Component: Terry Jernigan, Ph.D. (P.I.), Christine Fennema-Notestine, Ph.D., Sarah L. Archibald, M.A., John Hesselink, M.D., Jacopo Annese, Ph.D., Michael J. Taylor, Ph.D., Brian Schweinsburg, Ph.D.; Neurobiology Component: Eliezer Masliah, M.D. (P.I.), Ian P. Everall, FRCPsych., FRCPath., Ph.D., Cristian L. Achim, M.D., Ph.D.; Neurovirology Component: Douglas Richman, M.D., (P.I.), David M. Smith, M.D.; International Component: J. Allen McCutchan, M.D., (P.I.); Developmental Component: Ian P. Everall, FRCPsych., FRCPath., Ph.D. (P.I.), Stuart Lipton, M.D., Ph.D.;

Clinical Trials Component: J. Allen McCutchan, M.D., J. Hampton Atkinson, M.D., Ronald J. Ellis, M.D., Ph.D., Scott Letendre, M.D.; Participant Accrual and Retention Unit: J. Hampton Atkinson, M.D. (P.I.), Rodney von Jaeger, M.P.H.; Data Management Unit: Anthony C. Gamst, Ph.D. (P.I.), Clint Cushman (Data Systems Manager), Daniel R. Masys, M.D. (Senior Consultant); Statistics Unit: Ian Abramson, Ph.D. (P.I.), Florin Vaida, Ph.D., Christopher Ake, Ph.D.

C. L. Achim · I. P. Everall  
 Department of Psychiatry, University of California,  
 San Diego, CA, USA

C. L. Achim · W. Dumaop · E. Masliah  
 Department of Pathology, University of California,  
 San Diego, CA, USA

A. Adame · E. Masliah (✉)  
 Department of Neurosciences, University of California,  
 9500 Gilman Dr., La Jolla,  
 San Diego, CA 92093-0624, USA  
 e-mail: [emasliah@UCSD.edu](mailto:emasliah@UCSD.edu)

**Keywords** HIV · AIDS · amyloid · encephalitis · protein aggregation

## Introduction

In the age of highly active antiretroviral therapy (HAART), the survival of patients with AIDS has increased, and the proportion of HIV patients over the age of 50 has become significant. A recent study (Valcour et al. 2004a) found that individuals over 50 years of age represent 11% of the HIV cases registered by the Centers for Disease Control and Prevention (CDC). Data from the Hawaii cohort suggest an association between increased neurocognitive impairment in older HIV patients compared with younger controls (Valcour et al. 2004a, b, c, 2005; Valcour and Shiramizu 2004). Moreover, in a study in Western Pennsylvania (Becker et al. 2004), the prevalence of cognitive disorder among HIV-positive patients over 50 years of age was significantly higher than in younger individuals. Factors leading to this association may include vascular pathology, age-related immunological changes, and limited compensatory brain capacity. This suggests that chronic HIV infection and aging have similar effects on global cognitive functioning. Progressive brain dysfunction associated with aging has been linked to synapto-dendritic damage, reduced neurotransmitter production, and increased neuro-inflammatory reactivity associated with formation of radical oxygen species (Agrawal et al. 2006; Platenik et al. 2001; Price et al. 2006; Reynolds et al. 2007; Visalli et al. 2007).

Also contributing to the neuronal injury in older HIV patients might be the chronic permanence of HIV in the CNS. The HIV reservoir in the brain is known to be more difficult to reach because of the limited penetrance of HAART, emergence of resistant species, and viral escape (Dunfee et al. 2006; Langford et al. 2004).

Thus, despite the beneficial effects of HAART at suppressing viral load peripherally, the continued prevalence of moderate neurocognitive alterations in patients with HIV (Ances et al. 2008; Tozzi et al. 2007) might be related to the persistence of HIV in the CNS. Therefore, HIV encephalitis (HIVE) has shifted from a subacute neuro-inflammatory condition with abundant multinucleated giant cells to a more chronic and protracted condition with moderate astrogliosis, microgliosis, and viral load but more extensive neurodegeneration.

HAART has modified the patterns of HIV-related neuropathology and clinical manifestations in the past 10 years with the emergence of new variants of HIV encephalopathy (Everall et al. 2006). A paramount co-pathogenic factor in the long-term survival of patients with HIV on HAART is age. During aging, there is a failure in the handling and clearance of misfolded proteins such as A $\beta$  and  $\alpha$ -

synuclein. Misfolded proteins can be cleared via lysosomal (e.g., autophagy) and non-lysosomal pathways (e.g., proteasome; Williams et al. 2006). In Alzheimer's disease (AD) and other neurodegenerative disorders, this is manifested by increased accumulation of A $\beta$  into toxic oligomers (Barghorn et al. 2005; Deshpande et al. 2006; Glabe 2008; Shin et al. 2008; Standridge 2006; Tsigelny et al. 2008; Watson et al. 2005; White et al. 2005; Wisniewski and Konietzko 2008; Yoshiike et al. 2003; Selkoe 2008). Recent studies have suggested that in addition to the extracellular deposition of A $\beta$ , in AD, a smaller but significant fraction might accumulate intraneuronally (Arvanitis et al. 2007; Cuello 2005; Cuello and Canneva 2008; Echeverria and Cuello 2002; Grant et al. 2000; Green et al. 2005). Previous studies have shown that in aged HIV patients, there is increased amyloid deposition and plaque formation (Daily et al. 2006; Green et al. 2005; Rempel and Pulliam 2005). However, it was unclear what is the earliest site for A $\beta$  accumulation in patients with HIV and how this might be related to the alterations in protein handling during aging. For this purpose, patterns of intracellular A $\beta$  accumulation and its relationship with lysosomal clearance were analyzed in HIV cases ranging in age from 38–60 years old. Our results indicate that in this population, there is increased intraneuronal A $\beta$  accumulation in lysosomal-autophagic structures. These findings suggest that longer-term survival with HIV might interfere with clearance of proteins such as A $\beta$  that might worsen neuronal damage and cognitive impairment in this population.

## Materials and methods

**Subjects and neuropathological assessment** For the present study, we included a total of 43 HIV+ cases from the HIV Neurobehavioral Research Center and California Neuro-AIDS Tissue Network at the University of California San Diego. Cases had neuromedical and neuropsychological examinations within a median of 12 months before death. Most cases died as a result of acute bronchopneumonia or septicemia, and autopsy was performed within 24 h of death. Autopsy findings were consistent with AIDS, and the associated pathology was most frequently due to systemic CMV, Kaposi sarcoma, and liver disease. Subjects were excluded if they had a history of CNS opportunistic infections or non-HIV-related developmental, neurologic, psychiatric, or metabolic conditions that might affect CNS functioning (e.g., loss of consciousness exceeding 30 min, psychosis, substance dependence). In all cases, neuropathological assessment was performed in paraffin sections from the frontal, parietal, temporal cortices, hippocampus, basal ganglia, and brainstem stained with H&E or immunolabeled with antibodies against p24 and GFAP (Achim

et al. 1993; Masliah et al. 1992). The diagnosis of HIVE was based on the presence of microglial nodules, astrogliosis, HIV p24-positive cells, and myelin pallor. Additional analysis was performed with a subset of five age-matched (non-HIV) cases from the UCSD—Medical center Autopsy Service.

**Immunocytochemical analysis and image analysis** To further evaluate the patterns of A $\beta$  immunoreactivity in the brains of patients with HIV, briefly as previously described (Rockenstein et al. 2005) vibratome sections from the mid-frontal cortex (40  $\mu$ m thick) were incubated overnight at 4°C with the mouse monoclonal antibody against A $\beta$  (clone 4G8; 1:600, Senetek, Napa, CA, USA) or with the antibody against the N terminus of A $\beta$  (aa 1–16; clone 82E1; Immunobiological Laboratories, Gunma, Japan) followed by incubation with secondary biotinylated anti-mouse IgG, followed by ABC and DAB. Sections were transferred to SuperFrost slides (Fisher Scientific, Tustin, CA, USA) and mounted under glass coverslips with anti-fading media (Vector Laboratories, Burlingame, CA, USA). All sections were processed under the same standardized conditions. From each case, an average of 50 neurons were imaged, and digital files were analyzed with the Image Quant system. Three immunolabeled sections were analyzed per case, and the average of individual measurements was used to calculate group means. Additional confirmation of the intracellular amyloid labeling was performed utilizing the Thioflavine-S staining in the vibratome sections (Rockenstein et al. 2001).

**Double immunolabeling and confocal laser microscopy** To evaluate the co-localization between A $\beta$  immunoreactivity and lysosomal markers, double immunocytochemical analysis was performed (Belinson et al. 2008). For this purpose, vibratome sections were immunolabeled with a monoclonal antibody against A $\beta$  (clone 4G8; 1:600, Senetek, Napa, CA, USA) detected with FITC-conjugated secondary antibodies (1:75, Vector Laboratories) and the antibodies against cathepsin-D, Lamp2, and Rab5 (1:500, Dako, Carpinteria, CA, USA) detected with the Tyramide Signal Amplification™-Direct (Red) system (1:100, NEN Life Sciences, Boston, MA, USA; Pickford et al. 2008). All sections were processed simultaneously under the same

conditions, and experiments were performed twice to assess reproducibility. Sections were imaged with a Zeiss 63X (N.A. 1.4) objective on an Axiovert 35 microscope (Zeiss, Germany) with an attached MRC1024 LSCM system (BioRad). To confirm the specificity of primary antibodies, control experiments were performed where sections were incubated overnight in the absence of primary antibody (deleted) or preimmune serum and primary antibody alone.

**Immunoelectron microscopy analysis** Briefly, vibratome sections immunostained as described above with the 4G8 antibody or a nonimmune IgG were fixed in 0.25% glutaraldehyde and 3% paraformaldehyde in 0.1 M cacodylate buffer (pH 7.4) and then pre-embedded with 50% Durcupan epoxy resin, and 50% ethanol (dry) for 30 min. Samples were then embedded in Durcupan mix epoxy resin and polymerized under a vacuum at 60°C for 48 h. After the resin was polymerized, tissues were mounted into plastic cylinders, sectioned with an ultra microtome (Reichert Ultracut E) at 60 nm thickness and collected in copper grids for ultrastructural analysis. The immunostained grids were post-stained using saturated uranyl acetate solution in 50% ethanol for 20 min at room temperature, washed in distilled water, and placed in bismuth nitrate solution for 10 min followed by a final wash in deionized water. The immunolabeled grids were analyzed with a Zeiss EM10 electron microscope and electron micrographs obtained at a magnification of 35,000.


**Statistical analysis** All the analyses were conducted on blind-coded samples. After the results were obtained, the code was broken, and data were analyzed with the StatView program (SAS Institute, Cary, NC, USA). Comparisons among groups were performed with unpaired Student's *T* test, Chi square analysis and simple linear regression analysis. All results were expressed as mean $\pm$ SEM.

## Results

**Intraneuronal accumulation of A $\beta$  in HIV patients** A total of 48 cases were included, of which 43 were HIV seropositive, and five were HIV seronegative (Table 1). The age

**Table 1** Summary of demographic and pathological findings

Group	Number (N)	Gender M/F	Risk factor MM/DU/other	Age mean (years)	Age range (years)	Postmortem interval (h)	Cases with intracellular A $\beta$	Cases with amyloid plaques
HIV+, no HIVE	18	15/3	12/5/1	49 $\pm$ 1.5	38–60	18 $\pm$ 2	7/18 (38%)	1/18 (5%)
HIV+ Yes HIVE	25	21/4	16/9/0	47 $\pm$ 1	39–57	19 $\pm$ 2	18/25 (72%)	2/25 (8%)
HIV-	5	4/1	0/1/4	47 $\pm$ 2	38–55	15 $\pm$ 2	0/0	0/0




**Fig. 1** Patterns of A $\beta$  immunoreactivity in control and HIV+ cases. Panels are from the frontal cortex immunostained with the monoclonal antibody 4G8. **a–c** In an age-matched control HIV– case (42 year old) the neuronal cell bodies (**a**), axons in the white matter (**b**), and neuropil (**c**) are devoid of amyloid deposits. **d–f** Examples of intraneuronal (**d**) granular A $\beta$  immunoreactivity in an older HIV+ case (47 year old). A $\beta$  deposits can be found in axons (**e**) and in the neuropil (**f**) as diffuse plaques. **g–i** Examples of intraneuronal (**g**) and axonal (**h**) A $\beta$  immunoreactivity (*arrows*) in an older HIV+ case (50 year old) with HIV. The *inset* shows in greater detail the punctate appearance of the intraneuronal A $\beta$  immunostaining. Diffuse amyloid plaques (**i**) were also detected in a few cases. Bar=10  $\mu$ m


range varied between 38 and 60 years with a mean of  $48 \pm 2$  years. Of the 43 HIV cases, 18 had no significant opportunistic infections or HIV, and the other 25 had HIV. Immunocytochemical analysis with the antibody against A $\beta$  (4G8 clone) showed that compared to HIV– controls (Fig. 1A–C), in seven out of 18 HIV+ cases (38%) with no HIV, there was intraneuronal immunolabeling

(Fig. 1D). In contrast, in cases with HIV, intraneuronal A $\beta$  immunoreactivity was observed in 18 out of the 25 cases (72%; Fig. 1G). This difference was significant by Chi square analysis ( $X^2=4.7$ ,  $p=0.029$ ). The A $\beta$  immunostaining was observed in pyramidal neurons in layers 2–3 and 5, had a granular cytoplasmic appearance that was concentrated in the neuronal cell body (Figs. 1D, G; 2B–E), although in some cases, extended to the axons (Fig. 1E, H) and dendrites (Fig. 2D). In the five control cases, there was very low or no detectable intracellular A $\beta$  immunostaining (Figs. 1A–C, 2E). Image analysis of the levels of intracellular A $\beta$  immunostaining in the individual cases showed significantly higher levels in the HIV compared to HIV+ cases without HIV ( $t$  test,  $p=0.005$ ; Fig. 2). In one of the HIV+ cases with no apparent HIV (Fig. 1F) and in two of the cases with HIV, there was evidence of extracellular A $\beta$  deposition (Fig. 1I). The plaques had a diffuse appearance and in some cases surrounded neuronal cell bodies or were observed along axonal tracts (Fig. 3A–C). Although these diffuse plaques were similar to those observed in AD, no abundant neuritic plaques or tangles were detected, thus ruling out the possibility of AD in these cases. Similar results were observed with the antibody against the N terminus of A $\beta$  (82E1 clone) and with thioflavine-S (Fig. 3D–F). Linear regression analysis showed that there was a significant correlation between the levels of intracellular A $\beta$  immunoreactivity and age in the HIV+ group with HIV (Fig. 4A), but no correlation was observed in the HIV+ group with no HIV (Fig. 4B).

**Co-localization of lysosomal markers with the intraneuronal A $\beta$  in the brains of HIV patients** Given the punctate cytoplasmic characteristics of the intraneuronal A $\beta$  immunoreactivity in the HIV cases and that previous studies have suggested in experimental models that this might be localized



**Fig. 2** Levels of intraneuronal A $\beta$  immunoreactivity in older HIV+ cases. Images are from the frontal cortex immunostained with the monoclonal antibody 4G8. **a–d** Examples of the various levels (0–4) of intraneuronal A $\beta$  immunoreactivity in HIV+ cases; the *arrow* indicates




the enlarged image to the left displaying punctate appearance. **e** Compared to HIV+ with no HIV, in cases with HIV, there was an increase in the levels of intraneuronal A $\beta$  immunoreactivity. Bar=5  $\mu$ m



**Fig. 3** Laser confocal microscopy imaging of the amyloid deposits in HIV+ cases.

Examples are from the frontal cortex. **a** No evidence of amyloid deposits in HIV- age-matched control; **b, c** double-labeling with antibodies against the neuronal markers NeuN (red) and A $\beta$  (green) showing neuronal amyloid deposits (**b**, arrows) and diffuse plaques (**c**, amyl) in HIV+ cases. **d–f** Comparative images in HIV- (**d**) and HIV+ (**e, f**) cases stained with thioflavine S. Bar=10  $\mu$ m



in lysosomal structures, double labeling studies were performed. By confocal microscopy, the A $\beta$ -immunoreactive structures co-localized with the lysosomal markers cathepsin-D (Fig. 5D–F) and LAMP2 (Fig. 5 G–I). Moreover, the A $\beta$ -immunolabeled granular bodies co-localized with the autophagy marker LC3 (Fig. 5J–L). Overall, these granular structures were enlarged and clustered; in contrast, in control cases, discrete cathepsin-D lysosomal structures were identified that showed no A $\beta$  immunoreactivity (Fig. 5A–C).

To further investigate the intracellular localization of A $\beta$ , immunolabeled vibratome sections were analyzed by electron microscopy. Compared to control cases (Fig. 6A), in the HIV cases with intraneuronal A $\beta$  immunoreactivity, aggregates were observed associated with structures reminiscent of lysosomes and autophagolysosomes in the cell body and in axonal structures (Fig. 6B, C).


## Discussion

The present study showed that in HIV patients, there is a prominent intraneuronal accumulation of A $\beta$  associated

with lysosomal structures. The intraneuronal A $\beta$  was more abundant in patients with HIV compared to HIV cases with no significant pathology, and in the group with HIV, the levels of intracellular A $\beta$  increased with age. These findings suggest that long-term survival with HIV might interfere with clearance of proteins such as A $\beta$  that might worsen neuronal damage and cognitive impairment in this population. Previous studies have shown in older HIV+ patients the presence of extracellular amyloid deposits similar to those detected in patients with AD (Green et al. 2005; Rempel and Pulliam 2005).

The first study to identify changes similar to mild AD was reported by Esiri et al. (1998) who found A $\beta$  plaques in HIV patients by using argyrophilic and thioflavine stainings. Another study using a more comprehensive methodology that included staining with the 4G8 antibody, Congo red, and Thioflavine-S (Izycka-Swieszewska et al. 2000) showed that in 15 AIDS cases (five with HIV), three had perivascular plaques positive for 4G8 but not Congo Red or Thioflavine-S. Based on these findings, the authors concluded that the neurodegeneration associated with HIV infection could be primarily of vascular origin. Our results suggest that while

**Fig. 4** Linear regression analysis between intracellular A $\beta$  and age. **a** In cases with HIV, there was a significant correlation. **b** In cases with no HIV, there was no significant correlation





**Fig. 5** Laser confocal microscopy imaging of the intraneuronal A $\beta$  and lysosomal markers in older HIV+ cases.

Sections are from the frontal cortex double labeled with the A $\beta$  (4G8) antibody (*green*) and lysosomal proteins cathepsin-D, LAMP2, and LC3 (*red*).

**a–c** HIV- control case. **d–l** Co-localization (*arrows*) of intraneuronal A $\beta$  and cathepsin-D (**d–f**), LAMP2 (**g–i**), or LC3 (**j–l**) in an HIV+ case.

Bar=5  $\mu$ m



**Fig. 6** Ultrastructural analysis of the A $\beta$  deposits in HIV+ cases. Sections from the frontal cortex were immunostained pre-embedding with 4G8 and analyzed with the electron microscope. **a** In a control, HIV- case, evidence of preservation of neuronal organelles such as lysosomes (lys) and mitochondria (m). **b** In an HIV+ case, abnormal electrodense

membrane-bounded organelles (*arrows*) displaying A $\beta$  immunoreactivity accumulate in the neuronal cell body and axons. **c** Additional view of abnormal electrodense membrane-bounded organelles displaying intraneuronal A $\beta$  immunoreactivity in an HIV+ case. Bar=2  $\mu$ m

the vascular hypothesis may explain some of the plaques seen in the AIDS brains, given the intraneuronal accumulation and the neuronal origin of A $\beta$ , this supports a possible cerebral source of the parenchymal amyloid.

A $\beta$  is the result of the proteolytic processing of APP by  $\beta$ - and  $\gamma$ -secretases (Findeis 2007; Ghosh et al. 2005; Hoe and Rebeck 2008; Postina 2008; Russo et al. 2005; Thinakaran and Koo 2008; Vetrivel and Thinakaran 2006; Walsh et al. 2007; Wolfe 2006, 2008a, b). APP is ubiquitously distributed; in the CNS, APP is most abundantly produced by neurons and astroglial cells (Schmechel et al. 1988). Interestingly, the initial studies in the CNS of patients with HIV have focused on the levels of APP as a marker of neuronal injury. Several reports have described a significant increase in brain APP in AIDS cases, specifically in the axons of the subcortical white matter tracts. One leading theory is based on the inflammatory response in HIV infection of the brain parenchyma, where activated microglia are considered by many to be the likely source of mediators of disease that can promote overproduction and accumulation of APP (Adle-Biasette et al. 1999; Dickson et al. 1993). Among the brain macrophage-secreted factors mediating the overproduction of neuronal APP, a leading candidate is IL-1 (Stanley et al. 1994).

The association between the brain pathology characteristic of HIVE, including viral proteins and the presence of APP aggregates, often as intra-axonal globules, was reported in several studies (Giometto et al. 1997; Nebuloni et al. 2001; Raja et al. 1997). These results were further confirmed in the simian immunodeficiency virus (SIV) model of HIVE where Mankowski et al. have also found APP accumulation in degenerating axons (Mankowski et al. 2002). While the majority of these studies show a strong correlation between the topography of HIV-associated pathology and degenerating axons, suggesting a more local, intraparenchymal effect on APP accumulation, some investigators disagree. In one of the first studies of APP in the HIV brain, Scaravilli and colleagues (An et al. 1997) did not find a good correlation between APP and microgliosis and concluded that the axonal degeneration in HIV infection, may be due to systemic factors (e.g. cytokines).

The mechanisms through which HIV infection in the CNS might lead to increased accumulation of A $\beta$  are not completely clear. Recent studies suggest that HAART resistance, chronic HIV infection, HAART toxicity and reduced clearance of misfolded proteins might play a role (Hult et al. 2008). A $\beta$  is degraded by proteases such as neprilysin, insulin-degrading enzyme, and endothelin (Eckman and Eckman 2005; Miners et al. 2008; Nalivaeva et al. 2008; Wang et al. 2006). The HIV Tat protein has been shown to interfere with neprilysin activity, and HAART has been shown to alter the metabolic routes for A $\beta$  and insulin processing (Daily et al. 2006; Nath and

Hersh 2005; Rempel and Pulliam 2005). Neprilysin is not only capable of degrading extracellular but also intracellular A $\beta$  (Spencer et al. 2008). While the clinical benefit in reducing viral burden is indisputable, HAART is also reported to induce metabolic dysregulation resulting in a syndrome of lipodystrophy (LD) in up to 83% of treated individuals (Carr et al. 1999). Features associated with LD include insulin resistance with hyperinsulinemia, centripetal lipohypertrophy with concurrent subcutaneous peripheral lipodystrophy, and hypertriglyceridemia. The metabolic complications underlying LD include insulin resistance.

Another important mechanism involved in the clearance of A $\beta$  is the autophagy pathway (Jellinger 2006; Jellinger and Stadelmann 2001; Li et al. 2007; Lunemann et al. 2007; Nixon 2007; Shacka et al. 2008; Wilson et al. 2004; Yu et al. 2004, 2005; Zheng et al. 2006). This lysosomal pathway has been shown to be defective in patients with HIV (Delhay et al. 2007; Fredericksen et al. 2002; Levine and Sodora 2006; Tallozy et al. 2008), and recent studies in AD patients have shown that alterations in autophagy leads to increased intracellular and extracellular A $\beta$  (Pickford et al. 2008). Consistent with this possibility, in the present study, we showed the intracellular A $\beta$  co-localized with lysosomal markers and accumulated in structures reminiscent of lysosomes and abnormal autophagosomes.

In summary, in HIV patients, there is extensive intraneuronal accumulation of A $\beta$  suggesting that long-term survival with HIV might interfere with clearance of proteins such as A $\beta$  that might worsen neuronal damage and cognitive impairment in this population.

**Support** The study is supported by NIH grants MH076681, MH79881, MH45294, MH5974, MH58164, and DA12065, California NeuroAIDS Tissue Network U01 MH83506. The HIV Neurobehavioral Research Center (HNRC) is supported by Center award MH 62512 from NIMH.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

- Achim CL, Heyes MP, Wiley CA (1993) Quantitation of human immunodeficiency virus, immune activation factors, and quinolinic acid in AIDS brains. *J Clin Invest* 91:2769–2775 doi:10.1172/JCI116518
- Adle-Biasette H, Chretien F, Wingertsmann L, Hery C, Ereau T, Scaravilli F, Tardieu M, Gray F (1999) Neuronal apoptosis does not correlate with dementia in HIV infection but is related to microglial activation and axonal damage. *Neuropathol Appl Neurobiol* 25:123–133 doi:10.1046/j.1365-2990.1999.00167.x
- Agrawal L, Louboutin JP, Reyes BA, Van Bockstaele EJ, Strayer DS (2006) Antioxidant enzyme gene delivery to protect from HIV-1



- gp120-induced neuronal apoptosis. *Gene Ther* 13:1645–1656 doi:[10.1038/sj.gt.3302821](https://doi.org/10.1038/sj.gt.3302821)
- An SF, Giometto B, Groves M, Miller RF, Beckett AA, Gray F, Tavolato B, Scaravilli F (1997) Axonal damage revealed by accumulation of beta-APP in HIV-positive individuals without AIDS. *J Neuropathol Exp Neurol* 56:1262–1268
- Ances BM, Letendre SL, Alexander T, Ellis RJ (2008) Role of psychiatric medications as adjunct therapy in the treatment of HIV associated neurocognitive disorders. *Int Rev Psychiatry* 20:89–93 doi:[10.1080/09540260701877670](https://doi.org/10.1080/09540260701877670)
- Arvanitis DN, Ducatenzeiler A, Ou JN, Grodstein E, Andrews SD, Tendulkar SR, Ribeiro-da-Silva A, Szyf M, Cuello AC (2007) High intracellular concentrations of amyloid-beta block nuclear translocation of phosphorylated CREB. *J Neurochem* 103:216–228
- Barghorn S, Nimmrich V, Striebinger A, Krantz C, Keller P, Janson B, Bahr M, Schmidt M, Bitner RS, Harlan J, Barlow E, Ebert U, Hillen H (2005) Globular amyloid beta-peptide oligomer—a homogenous and stable neuropathological protein in Alzheimer's disease. *J Neurochem* 95:834–847 doi:[10.1111/j.1471-4159.2005.03407.x](https://doi.org/10.1111/j.1471-4159.2005.03407.x)
- Becker JT, Lopez OL, Dew MA, Aizenstein HJ (2004) Prevalence of cognitive disorders differs as a function of age in HIV virus infection. *AIDS* 18(Suppl 1):S11–S18 doi:[10.1097/00002030-200401001-00003](https://doi.org/10.1097/00002030-200401001-00003)
- Belinson H, Lev D, Masliah E, Michaelson DM (2008) Activation of the amyloid cascade in apolipoprotein E4 transgenic mice induces lysosomal activation and neurodegeneration resulting in marked cognitive deficits. *J Neurosci* 28:4690–4701 doi:[10.1523/JNEUROSCI.5633-07.2008](https://doi.org/10.1523/JNEUROSCI.5633-07.2008)
- Carr A, Samaras K, Thorisdottir A, Kaufmann GR, Chisholm DJ, Cooper DA (1999) Diagnosis, prediction, and natural course of HIV-1 protease-inhibitor-associated lipodystrophy, hyperlipidaemia, and diabetes mellitus: a cohort study. *Lancet* 353:2093–2099 doi:[10.1016/S0140-6736\(98\)08468-2](https://doi.org/10.1016/S0140-6736(98)08468-2)
- Cuello AC (2005) Intracellular and extracellular Abeta, a tale of two neuropathologies. *Brain Pathol* 15:66–71
- Cuello AC, Caneva F (2008) Impact of intracellular beta-amyloid in transgenic animals and cell models. *Neurodegener Dis* 5:146–148 doi:[10.1159/000113686](https://doi.org/10.1159/000113686)
- Daily A, Nath A, Hersh LB (2006) Tat peptides inhibit neprilysin. *J Neurovirol* 12:153–160 doi:[10.1080/13550280600760677](https://doi.org/10.1080/13550280600760677)
- Delhay M, Gravot A, Ayinde D, Niedergang F, Alizon M, Brelot A (2007) Identification of a postendocytic sorting sequence in CCR5. *Mol Pharmacol* 72:1497–1507 doi:[10.1124/mol.107.038422](https://doi.org/10.1124/mol.107.038422)
- Deshpande A, Mina E, Glabe C, Busciglio J (2006) Different conformations of amyloid beta induce neurotoxicity by distinct mechanisms in human cortical neurons. *J Neurosci* 26:6011–6018 doi:[10.1523/JNEUROSCI.1189-06.2006](https://doi.org/10.1523/JNEUROSCI.1189-06.2006)
- Dickson DW, Lee SC, Mattiace LA, Yen SH, Brosnan C (1993) Microglia and cytokines in neurological disease, with special reference to AIDS and Alzheimer's disease. *Glia* 7:75–83 doi:[10.1002/glia.440070113](https://doi.org/10.1002/glia.440070113)
- Dunfee R, Thomas ER, Gorry PR, Wang J, Ancuta P, Gabuzda D (2006) Mechanisms of HIV-1 neurotropism. *Curr HIV Res* 4:267–278 doi:[10.2174/15701620677709500](https://doi.org/10.2174/15701620677709500)
- Echeverria V, Cuello AC (2002) Intracellular A-beta amyloid, a sign for worse things to come? *Mol Neurobiol* 26:299–316 doi:[10.1385/MN:26:2-3:299](https://doi.org/10.1385/MN:26:2-3:299)
- Eckman EA, Eckman CB (2005) Abeta-degrading enzymes: modulators of Alzheimer's disease pathogenesis and targets for therapeutic intervention. *Biochem Soc Trans* 33:1101–1105 doi:[10.1042/BST20051101](https://doi.org/10.1042/BST20051101)
- Esiri MM, Biddolph SC, Morris CS (1998) Prevalence of Alzheimer plaques in AIDS. *J Neurol Neurosurg Psychiatry* 65:29–33 doi:[10.1136/jnnp.65.1.29](https://doi.org/10.1136/jnnp.65.1.29)
- Everall IP, Salaria S, Atkinson JH, Young C, Corbeil J, Grant I, Masliah E (2006) Diminished somatostatin gene expression in individuals with HIV and major depressive disorder. *Neurology* 67:1867–1869 doi:[10.1212/01.wnl.0000244436.04036.a2](https://doi.org/10.1212/01.wnl.0000244436.04036.a2)
- Findeis MA (2007) The role of amyloid beta peptide 42 in Alzheimer's disease. *Pharmacol Ther* 116:266–286 doi:[10.1016/j.pharmthera.2007.06.006](https://doi.org/10.1016/j.pharmthera.2007.06.006)
- Fredericksen BL, Wei BL, Yao J, Luo T, Garcia JV (2002) Inhibition of endosomal/lysosomal degradation increases the infectivity of human immunodeficiency virus. *J Virol* 76:11440–11446 doi:[10.1128/JVI.76.22.11440-11446.2002](https://doi.org/10.1128/JVI.76.22.11440-11446.2002)
- Ghosh AK, Kumaragurubaran N, Tang J (2005) Recent developments of structure based beta-secretase inhibitors for Alzheimer's disease. *Curr Top Med Chem* 5:1609–1622 doi:[10.2174/156802605775009711](https://doi.org/10.2174/156802605775009711)
- Giometto B, An SF, Groves M, Scaravilli T, Geddes JF, Miller R, Tavolato B, Beckett AA, Scaravilli F (1997) Accumulation of beta-amyloid precursor protein in HIV encephalitis: relationship with neuropsychological abnormalities. *Ann Neurol* 42:34–40 doi:[10.1002/ana.410420108](https://doi.org/10.1002/ana.410420108)
- Glabe CG (2008) Structural classification of toxic amyloid oligomers. *J Biol Chem* 283:29639–29643 doi:[10.1074/jbc.R800016200](https://doi.org/10.1074/jbc.R800016200)
- Grant SM, Ducatenzeiler A, Szyf M, Cuello AC (2000) Abeta immunoreactive material is present in several intracellular compartments in transfected, neuronally differentiated, P19 cells expressing the human amyloid beta-protein precursor. *J Alzheimers Dis* 2:207–222
- Green DA, Masliah E, Vinters HV, Beizai P, Moore DJ, Achim CL (2005) Brain deposition of beta-amyloid is a common pathologic feature in HIV positive patients. *AIDS* 19:407–411 doi:[10.1097/01.aids.0000161770.06158.5c](https://doi.org/10.1097/01.aids.0000161770.06158.5c)
- Hoe HS, Rebeck GW (2008) Regulated proteolysis of APP and ApoE receptors. *Mol Neurobiol* 37:64–72 doi:[10.1007/s12035-008-8017-0](https://doi.org/10.1007/s12035-008-8017-0)
- Hult B, Chana G, Masliah E, Everall I (2008) Neurobiology of HIV. *Int Rev Psychiatry* 20:3–13 doi:[10.1080/09540260701862086](https://doi.org/10.1080/09540260701862086)
- Izycka-Swieszewska E, Zoltowska A, Rzepko R, Gross M, Borowska-Lehman J (2000) Vasculopathy and amyloid beta reactivity in brains of patients with acquired immune deficiency (AIDS). *Folia Neuropathol* 38:175–182
- Jellinger KA (2006) Challenges in neuronal apoptosis. *Curr Alzheimer Res* 3:377–391 doi:[10.2174/156720506778249434](https://doi.org/10.2174/156720506778249434)
- Jellinger KA, Stadelmann C (2001) Problems of cell death in neurodegeneration and Alzheimer's Disease. *J Alzheimers Dis* 3:31–40
- Langford D, Grigorian A, Hurford R, Adame A, Ellis RJ, Hansen L, Masliah E (2004) Altered P-glycoprotein expression in AIDS patients with HIV encephalitis. *J Neuropathol Exp Neurol* 63:1038–1047
- Levine B, Sodora DL (2006) HIV and CXCR4 in a kiss of autophagic death. *J Clin Invest* 116:2078–2080 doi:[10.1172/JCI29447](https://doi.org/10.1172/JCI29447)
- Li M, Chen L, Lee DH, Yu LC, Zhang Y (2007) The role of intracellular amyloid beta in Alzheimer's disease. *Prog Neurobiol* 83:131–139 doi:[10.1016/j.pneurobio.2007.08.002](https://doi.org/10.1016/j.pneurobio.2007.08.002)
- Lunemann JD, Schmidt J, Schmid D, Barthel K, Wrede A, Dalakas MC, Munz C (2007) Beta-amyloid is a substrate of autophagy in sporadic inclusion body myositis. *Ann Neurol* 61:476–483 doi:[10.1002/ana.21115](https://doi.org/10.1002/ana.21115)
- Mankowski JL, Queen SE, Tarwater PM, Fox KJ, Perry VH (2002) Accumulation of beta-amyloid precursor protein in axons correlates with CNS expression of SIV gp41. *J Neuropathol Exp Neurol* 61:85–90
- Masliah E, Achim CL, Ge N, DeTeresa R, Terry RD, Wiley CA (1992) Spectrum of human immunodeficiency virus-associated neocortical damage. *Ann Neurol* 32:321–329 doi:[10.1002/ana.410320304](https://doi.org/10.1002/ana.410320304)
- Miners JS, Baig S, Palmer J, Palmer LE, Kehoe PG, Love S (2008) Abeta-degrading enzymes in Alzheimer's disease. *Brain Pathol* 18:240–252 doi:[10.1111/j.1750-3639.2008.00132.x](https://doi.org/10.1111/j.1750-3639.2008.00132.x)



- Nalivaeva NN, Fisk LR, Belyaev ND, Turner AJ (2008) Amyloid-degrading enzymes as therapeutic targets in Alzheimer's disease. *Curr Alzheimer Res* 5:212–224 doi:[10.2174/156720508783954785](https://doi.org/10.2174/156720508783954785)
- Nath A, Hersh LB (2005) Tat and amyloid: multiple interactions. *AIDS* 19:203–204 doi:[10.1097/00002030-200501280-00013](https://doi.org/10.1097/00002030-200501280-00013)
- Nebuloni M, Pellegrinelli A, Ferri A, Bonetto S, Boldorini R, Vago L, Grassi MP, Costanzi G (2001) Beta amyloid precursor protein and patterns of HIV p24 immunohistochemistry in different brain areas of AIDS patients. *AIDS* 15:571–575 doi:[10.1097/00002030-200103300-00005](https://doi.org/10.1097/00002030-200103300-00005)
- Nixon RA (2007) Autophagy, amyloidogenesis and Alzheimer disease. *J Cell Sci* 120:4081–4091 doi:[10.1242/jcs.019265](https://doi.org/10.1242/jcs.019265)
- Pickford F, Masliah E, Britschgi M, Lucin K, Narasimhan R, Jaeger PA, Small S, Spencer B, Rockenstein E, Levine B, Wyss-Coray T (2008) The autophagy-related protein beclin 1 shows reduced expression in early Alzheimer disease and regulates amyloid beta accumulation in mice. *J Clin Invest* 118:2190–2199
- Platenik J, Stopka P, Vejrazka M, Stipek S (2001) Quinolinic acid-iron (ii) complexes: slow autooxidation, but enhanced hydroxyl radical production in the Fenton reaction. *Free Radic Res* 34:445–459 doi:[10.1080/10715760100300391](https://doi.org/10.1080/10715760100300391)
- Postina R (2008) A closer look at alpha-secretase. *Curr Alzheimer Res* 5:179–186 doi:[10.2174/156720508783954668](https://doi.org/10.2174/156720508783954668)
- Price TO, Uras F, Banks WA, Ercal N (2006) A novel antioxidant *N*-acetylcysteine amide prevents gp120- and Tat-induced oxidative stress in brain endothelial cells. *Exp Neurol* 201:193–202 doi:[10.1016/j.expneurol.2006.03.030](https://doi.org/10.1016/j.expneurol.2006.03.030)
- Raja F, Sherriff FE, Morris CS, Bridges LR, Esiri MM (1997) Cerebral white matter damage in HIV infection demonstrated using beta-amyloid precursor protein immunoreactivity. *Acta Neuropathol* 93:184–189 doi:[10.1007/s004010050601](https://doi.org/10.1007/s004010050601)
- Rempel HC, Pulliam L (2005) HIV-1 Tat inhibits neprilysin and elevates amyloid beta. *AIDS* 19:127–135 doi:[10.1097/00002030-200501280-00004](https://doi.org/10.1097/00002030-200501280-00004)
- Reynolds A, Laurie C, Mosley RL, Gendelman HE (2007) Oxidative stress and the pathogenesis of neurodegenerative disorders. *Int Rev Neurobiol* 82:297–325 doi:[10.1016/S0074-7742\(07\)82016-2](https://doi.org/10.1016/S0074-7742(07)82016-2)
- Rockenstein E, Mallory M, Mante M, Sisk A, Masliah E (2001) Early formation of mature amyloid-beta protein deposits in a mutant APP transgenic model depends on levels of Aβ(1–42). *J Neurosci Res* 66:573–582 doi:[10.1002/jnr.1247](https://doi.org/10.1002/jnr.1247)
- Rockenstein E, Schwach G, Ingolic E, Adame A, Crews L, Mante M, Pfragner R, Schreiner E, Windisch M, Masliah E (2005) Lysosomal pathology associated with alpha-synuclein accumulation in transgenic models using an eGFP fusion protein. *J Neurosci Res* 80:247–259 doi:[10.1002/jnr.20446](https://doi.org/10.1002/jnr.20446)
- Russo C, Venezia V, Repetto E, Nizzari M, Violani E, Carlo P, Schettini G (2005) The amyloid precursor protein and its network of interacting proteins: physiological and pathological implications. *Brain Res Brain Res Rev* 48:257–264 doi:[10.1016/j.brainresrev.2004.12.016](https://doi.org/10.1016/j.brainresrev.2004.12.016)
- Schmechel DE, Goldgaber D, Burkhart DS, Gilbert JR, Gajdusek DC, Roses AD (1988) Cellular localization of messenger RNA encoding amyloid-beta-protein in normal tissue and in Alzheimer disease. *Alzheimer Dis Assoc Disord* 2:96–111 doi:[10.1097/00002093-198802020-00002](https://doi.org/10.1097/00002093-198802020-00002)
- Selkoe DJ (2008) Soluble oligomers of the amyloid beta-protein impair synaptic plasticity and behavior. *Behav Brain Res* 192:106–113 doi:[10.1016/j.bbr.2008.02.016](https://doi.org/10.1016/j.bbr.2008.02.016)
- Shacka JJ, Roth KA, Zhang J (2008) The autophagy-lysosomal degradation pathway: role in neurodegenerative disease and therapy. *Front Biosci* 13:718–736 doi:[10.2741/2714](https://doi.org/10.2741/2714)
- Shin TM, Isas JM, Hsieh CL, Kaye R, Glabe CG, Langen R, Chen J (2008) Formation of soluble amyloid oligomers and amyloid fibrils by the multifunctional protein vitronectin. *Mol Neurodegener* 3:16 doi:[10.1186/1750-1326-3-16](https://doi.org/10.1186/1750-1326-3-16)
- Spencer B, Marr RA, Rockenstein E, Crews L, Adame A, Potkar R, Patrick C, Gage FH, Verma IM, Masliah E (2008) Long-term neprilysin gene transfer is associated with reduced levels of intracellular Aβ and behavioral improvement in APP transgenic mice. *BMC Neurosci* 9:109 doi:[10.1186/1471-2202-9-109](https://doi.org/10.1186/1471-2202-9-109)
- Standridge JB (2006) Vicious cycles within the neuropathophysiologic mechanisms of Alzheimer's disease. *Curr Alzheimer Res* 3:95–108 doi:[10.2174/156720506776383068](https://doi.org/10.2174/156720506776383068)
- Stanley LC, Mrak RE, Woody RC, Perrot LJ, Zhang S, Marshak DR, Nelson SJ, Griffin WS (1994) Glial cytokines as neuropathogenic factors in HIV infection: pathogenic similarities to Alzheimer's disease. *J Neuropathol Exp Neurol* 53:231–238 doi:[10.1097/00005072-199405000-00003](https://doi.org/10.1097/00005072-199405000-00003)
- Tallosy Z, Martinez J, Joset D, Ray Y, Gacser A, Toussi S, Mizushima N, Nosanchuk J, Goldstein H, Loike J, Sulzer D, Santambrogio L (2008) Methamphetamine inhibits antigen processing, presentation, and phagocytosis. *PLoS Pathog* 4:e28 doi:[10.1371/journal.ppat.0040028](https://doi.org/10.1371/journal.ppat.0040028)
- Thinakaran G, Koo EH (2008) Amyloid precursor protein trafficking, processing, and function. *J Biol Chem* 283:29615–29619 doi:[10.1074/jbc.R800019200](https://doi.org/10.1074/jbc.R800019200)
- Tozzi V, Balestra P, Bellagamba R, Corpolongo A, Salvatori MF, Visco-Comandini U, Vlasi C, Giulianelli M, Galgani S, Antinori A, Narciso P (2007) Persistence of neuropsychologic deficits despite long-term highly active antiretroviral therapy in patients with HIV-related neurocognitive impairment: prevalence and risk factors. *J Acquir Immune Defic Syndr* 45:174–182 doi:[10.1097/QAI.0b013e318042e1ee](https://doi.org/10.1097/QAI.0b013e318042e1ee)
- Tsigelny IF, Crews L, Desplats P, Shaked GM, Sharikov Y, Mizuno H, Spencer B, Rockenstein E, Trejo M, Platoshyn O, Yuan JX, Masliah E (2008) Mechanisms of hybrid oligomer formation in the pathogenesis of combined Alzheimer's and Parkinson's diseases. *PLoS One* 3:e3135 doi:[10.1371/journal.pone.0003135](https://doi.org/10.1371/journal.pone.0003135)
- Valcour V, Shiramizu B (2004) HIV-associated dementia, mitochondrial dysfunction, and oxidative stress. *Mitochondrion* 4:119–129 doi:[10.1016/j.mito.2004.05.009](https://doi.org/10.1016/j.mito.2004.05.009)
- Valcour V, Shikuma C, Shiramizu B, Watters M, Poff P, Selnes O, Holck P, Grove J, Sacktor N (2004a) Higher frequency of dementia in older HIV-1 individuals: the Hawaii Aging with HIV-1 Cohort. *Neurology* 63:822–827
- Valcour V, Shikuma C, Shiramizu B, Watters M, Poff P, Selnes OA, Grove J, Liu Y, Abdul-Majid KB, Gartner S, Sacktor N (2004b) Age, apolipoprotein E4, and the risk of HIV dementia: the Hawaii Aging with HIV Cohort. *J Neuroimmunol* 157:197–202 doi:[10.1016/j.jneuroim.2004.08.029](https://doi.org/10.1016/j.jneuroim.2004.08.029)
- Valcour VG, Shikuma CM, Watters MR, Sacktor NC (2004c) Cognitive impairment in older HIV-1-seropositive individuals: prevalence and potential mechanisms. *AIDS* 18(Suppl 1):S79–S86 doi:[10.1097/00002030-200401001-00012](https://doi.org/10.1097/00002030-200401001-00012)
- Valcour VG, Shikuma CM, Shiramizu BT, Williams AE, Watters MR, Poff PW, Grove JS, Selnes OA, Sacktor NC (2005) Diabetes, insulin resistance, and dementia among HIV-1-infected patients. *J Acquir Immune Defic Syndr* 38:31–36 doi:[10.1097/00126334-200501010-00006](https://doi.org/10.1097/00126334-200501010-00006)
- Vetrivel KS, Thinakaran G (2006) Amyloidogenic processing of beta-amyloid precursor protein in intracellular compartments. *Neurology* 66:S69–S73 doi:[10.1212/01.wnl.0000192107.17175.39](https://doi.org/10.1212/01.wnl.0000192107.17175.39)
- Visalli V, Muscoli C, Sacco I, Sculco F, Palma E, Costa N, Colica C, Rotiroti D, Mollace V (2007) *N*-acetylcysteine prevents HIV gp 120-related damage of human cultured astrocytes: correlation with glutamine synthase dysfunction. *BMC Neurosci* 8:106 doi:[10.1186/1471-2202-8-106](https://doi.org/10.1186/1471-2202-8-106)
- Walsh DM, Minogue AM, Sala Frigerio C, Fadeeva JV, Wasco W, Selkoe DJ (2007) The APP family of proteins: similarities and differences. *Biochem Soc Trans* 35:416–420 doi:[10.1042/BST0350416](https://doi.org/10.1042/BST0350416)

- Wang DS, Dickson DW, Malter JS (2006) Beta-amyloid degradation and Alzheimer's disease. *J Biomed Biotechnol* 2006:58406 doi:[10.1155/JBB/2006/58406](https://doi.org/10.1155/JBB/2006/58406)
- Watson D, Castano E, Kokjohn TA, Kuo YM, Lyubchenko Y, Pinsky D, Connolly ES Jr, Esh C, Luehrs DC, Stine WB, Rowse LM, Emmerling MR, Roher AE (2005) Physicochemical characteristics of soluble oligomeric A $\beta$  and their pathologic role in Alzheimer's disease. *Neurol Res* 27:869–881 doi:[10.1179/016164105X49436](https://doi.org/10.1179/016164105X49436)
- White JA, Manelli AM, Holmberg KH, Van Eldik LJ, Ladu MJ (2005) Differential effects of oligomeric and fibrillar amyloid-beta 1–42 on astrocyte-mediated inflammation. *Neurobiol Dis* 18:459–465 doi:[10.1016/j.nbd.2004.12.013](https://doi.org/10.1016/j.nbd.2004.12.013)
- Williams A, Jahreiss L, Sarkar S, Saiki S, Menzies FM, Ravikumar B, Rubinsztein DC (2006) Aggregate-prone proteins are cleared from the cytosol by autophagy: therapeutic implications. *Curr Top Dev Biol* 76:89–101 doi:[10.1016/S0070-2153\(06\)76003-3](https://doi.org/10.1016/S0070-2153(06)76003-3)
- Wilson CA, Murphy DD, Giasson BI, Zhang B, Trojanowski JQ, Lee VM (2004) Degradative organelles containing mislocalized alpha-and beta-synuclein proliferate in presenilin-1 null neurons. *J Cell Biol* 165:335–346 doi:[10.1083/jcb.200403061](https://doi.org/10.1083/jcb.200403061)
- Wisniewski T, Konietzko U (2008) Amyloid-beta immunisation for Alzheimer's disease. *Lancet Neurol* 7:805–811 doi:[10.1016/S1474-4422\(08\)70170-4](https://doi.org/10.1016/S1474-4422(08)70170-4)
- Wolfe MS (2006) The gamma-secretase complex: membrane-embedded proteolytic ensemble. *Biochemistry* 45:7931–7939 doi:[10.1021/bi060799c](https://doi.org/10.1021/bi060799c)
- Wolfe MS (2008a) Gamma-secretase inhibition and modulation for Alzheimer's disease. *Curr Alzheimer Res* 5:158–164 doi:[10.2174/156720508783954767](https://doi.org/10.2174/156720508783954767)
- Wolfe MS (2008b) Gamma-secretase: structure, function, and modulation for Alzheimer's disease. *Curr Top Med Chem* 8:2–8 doi:[10.2174/156802608783334024](https://doi.org/10.2174/156802608783334024)
- Yoshiike Y, Chui DH, Akagi T, Tanaka N, Takashima A (2003) Specific compositions of amyloid-beta peptides as the determinant of toxic beta-aggregation. *J Biol Chem* 278:23648–23655 doi:[10.1074/jbc.M212785200](https://doi.org/10.1074/jbc.M212785200)
- Yu WH, Kumar A, Peterhoff C, Shapiro Kulnane L, Uchiyama Y, Lamb BT, Cuervo AM, Nixon RA (2004) Autophagic vacuoles are enriched in amyloid precursor protein-secretase activities: implications for beta-amyloid peptide over-production and localization in Alzheimer's disease. *Int J Biochem Cell Biol* 36:2531–2540 doi:[10.1016/j.biocel.2004.05.010](https://doi.org/10.1016/j.biocel.2004.05.010)
- Yu WH, Cuervo AM, Kumar A, Peterhoff CM, Schmidt SD, Lee JH, Mohan PS, Mercken M, Farmery MR, Tjernberg LO, Jiang Y, Duff K, Uchiyama Y, Naslund J, Mathews PM, Cataldo AM, Nixon RA (2005) Macroautophagy—a novel Beta-amyloid peptide-generating pathway activated in Alzheimer's disease. *J Cell Biol* 171:87–98 doi:[10.1083/jcb.200505082](https://doi.org/10.1083/jcb.200505082)
- Zheng L, Roberg K, Jerhammar F, Marcusson J, Terman A (2006) Oxidative stress induces intralysosomal accumulation of Alzheimer amyloid beta-protein in cultured neuroblastoma cells. *Ann N Y Acad Sci* 1067:248–251 doi:[10.1196/annals.1354.032](https://doi.org/10.1196/annals.1354.032)